PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:

A61K 9/20, 9/50, 9/52 A61K 31/19

(11) International Publication Number:

WO 91/15194

A1

(43) International Publication Date:

17 October 1991 (17.10.91)

(21) International Application Number:

PCT/US91/01089

(22) International Filing Date:

26 February 1991 (26.02.91)

(30) Priority data:

508,193

11 April 1990 (11.04.90) US

(60) Parent Application or Grant

(63) Related by Continuation US

508,193 (CON)

Filed on

11 April 1990 (11.04.90)

(71) Applicant (for all designated States except US): THE UP-JOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): SHEN, Robert, Wu-Wei [US/US]; 5121 Maple Ridge Road, Kalamazoo, MI 49008 (US).

(74) Agent: WILLIAMS, Sidney, B., Jr.; Corporate Patents & Trademarks, The Upjohn Company, Kalamazoo, MI 49001 (US).

(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent), DK (European patent), ES (EUROPEAN EUROPEAN EUROPEA tent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW NL (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPÍ patent), US.

Published

With international search report.

(54) Title: TASTE MASKING OF IBUPROFEN BY FLUID BED COATING

(57) Abstract

A chewable taste-masked ibuprofen tablet having controlled release characteristics.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

			•		
AT	Austria	ES	Spain	MG	Madagascar
ΑU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	
BJ	Benin	GR	Greece		Netherlands
BR	Brazil	HU		NO	Norway
CA	Canada	_	Hungary	PL	Poland
CF	=	ΙΤ	Italy	RO	Romania
-	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic	SE	Sweden
CH	Switzerland		of Korca	SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
cs	Czechoslovakia	LK	Sri Lanka	TG	
DE	Germany	LU	Luxembourg		Togo
DK	Denmark	MC	Monaco	US	United States of America

15

20

25

TASTE MASKING OF IBUPROFEN BY FLUID BED COATING BACKGROUND OF THE INVENTION

Ibuprofen is a well-known therapeutic agent. Its therapeutic activities include analgesia and anti-pyretic attack. As with most medicines, one of the difficulties with ibuprofen is in making it palatable to children. This difficulty has been overcome with most medicines by preparing formulations such as syrups and drops. The present invention relates to chewable tablets that are palatable to children and a process for making the tablets.

From a manufacturing cost standpoint, it is desirable to have chewable, taste-masked microcapsules that are large (0.25-1 mm in diameter), because larger microcapsules are easier to manufacture and package, and are less expensive to produce than are small microcapsules. However, an increase in size makes fracture during chewing and the release of drug from the microcapsule more likely to occur especially when there is an inadequate amount of plasticizer or other component included to provide elasticity. A larger sized microcapsule requires greater elasticity to minimize the likelihood that a fracture will occur and active agent will be released. There is therefore a need in the art of pharmaceutical formulation to provide encapsulating coatings capable of being formulated into chewable microcapsules as large as about 1.5 mm. that will not release drugs during chewing.

INFORMATION DISCLOSURE

Ibuprofen and its use for treatment of analgesia is disclosed in U.S. patent 3,385,886. Compositions containing ibuprofen and methods for using them are described in U.S. patent 3,228,831. New crystalline and high dose formulations of ibuprofen are disclosed in U.S. patents 4,476,248 and 4,609,675 respectively.

Microencapsulation is described by J.A. Bakan, Part Three of "The Theory and Practice of Industrial Pharmacy", 1986, pp. 413-429.

EUDRAGIT L30D is a known polymer useful for coating orally administered pharmaceutical dosage forms, particularly tablets, capsules and pills, with coatings which are resistant to gastric juices but solvent in intestinal juices.

30 Chewable taste-masked pharmaceutical compositions, including some containing ibuprofen, are described in U.S. patent 4,800,087. However, the compositions described therein require a coating consisting of a mixture of polymers. The use of fluidized bed for coating pharmaceutical products is described in U.S. Patent 4,800,087.

10

15

20

-2-

SUMMARY OF THE INVENTION

This invention involves:

A chewable taste-masked tablet having controlled release characteristics comprising a microcapsule of about 100 microns to about 0.8 mm in diameter having (a) a pharmaceutical core including crystalline ibuprofen and (b) a methacrylic acid copolymer coating having sufficient elasticity to withstand chewing.

While chewable taste masked formulations of ibuprofen are referred to in the prior art, among the advantages of the compositions of this invention over the closest prior art compositions is that the coating used consists of a single copolymer rather than a mixture of copolymers.

DETAILED DESCRIPTION OF THE INVENTION

The present invention comprises formulations of taste-masked microcapsules which further comprise (a) a pharmaceutical core of crystalline ibuprofen and (b) a methacrylic acid copolymer coating that may provide chewable taste-masked characteristics. Both the polymeric coating and the pharmaceutical core may further comprise diluents, fillers and other pharmaceutical additives which may effect the rate of release of active ibuprofen from the microcapsule.

The methacrylic acid copolymer is preferably dispersable in water so as to take advantage of aqueous formulation techniques and has a rapid rate of dissolution at a pH of about 5.5.. Aqueous-based coating systems are safe and make regulatory compliance (EPA) relatively easy compared to non-aquaous based coating systems. An elastic microcapsule which will not release ibuprofen in the mouth when chewed is contemplated by the present invention.

A preferred coating composition is a high temperature film forming polymer or 25 "hard" polymer. A hard polymer is defined as a polymer that will form a film on a pharmaceutical core at a temperature of at least about 30°C. Examples of high temperature film forming polymers useful in this invention include hydroxypropylmethyl cellulose, for example, Pharmacoat' 606 brand from Shinetro Corp., Tokyo, Japan, hydroxypropyl cellulose, for example, Klucel' brand from Hercules Corp., Wilmington, Del., methylcellulose for example Methocel A', from Dow Chemical, Midland, Mich., ethylcellulose, for example, Ethocel' brand from Dow Chemical Corp., and other aqueous polymeric dispersions such as Aquacost' Brand from FMC, Philadelphia, PA., and Surelease' brand from Colorcon, West Point, PA., polyvinyl alcohol, polyvinyl

10

15

20

25

30

acetate, cellulose acetate butyrate, styrene acylate copolymers, for example Janocryl 138 (61°C. film forming copolymer from S.C. Johnson, Racine, Wis.) and copolymers of acrylic acid esters, for example, the EUDRAGIT Copolymers (Rohm Pharma GmbH) Westerstadt, W. Germany): Eudragit L30D, Eudragit L100-55, Eudragit RS(30D and 100).

Eudragit' copolymers that are preferred in embodiments of this invention include L30D, an anionic copolymer based of polymethacrylic and acrylic acid esters (Methacrylic Acid Copolymer, Type C in USP XXL/NF XVI) with a mean molecular weight of 250,000.

The polymeric coating should provide for immediate release characteristics, i.e., rapid release of the active agents in the duodenum within a period of about one hour. When the microcapsules are formulated into chewable, taste-masked oral tablets or capsules, the formulations provide for immediate, rapid release in the stomach.

The chewable polymeric coating providing immediate release upon reaching the duodenum i.e., within one hour after ingestion may be comprised of a pharmaceutically compatible high temperature film forming polymer that is water insoluble or not swellable within the pH range (about 5.5-6.5) and/or the liquid content of the mouth and will not release the active agent in the mouth, but will dissolve or change in physical character in the duodenum, for example, swell or become more porous, thus releasing drug.

The most preferred film forming acrylic resin polymer that releases active agent rapidly in duodenum is EUDRAGIT L30D. EUDRAGIT L30D is a copolymer anionic in character, based on polymethacrylic acid and acrylic acid esters. Although EUDRAGIT L30D is soluble at pH's in the mouth and insoluble at pH's of the stomach, it has found usefulness in chewable, taste-masked immediate release formulations of the present invention. This usefulness may stem from the lack of liquid in the mouth, or may be the result of elastic qualities that EUDRAGIT L30D acquires when formulated in combination with a plasticizer, or preferably, with EUDRAGIT E30D.

Any of the above described high-temperature film-forming polymers may be used for microencapsulation. However, to make capsules of the required elasticity using the above described high temperature film forming polymers, plasticizers may be incorporated into the coatings. Plasticizers useful to provide the requisite elasticity include propylene glycol and polyalkylene glycols, for example, polyethylene glycol;

triacetin, (glyceryl triacetate from Eastman Kodak, Rochester, NY vinyl pyrrolidone, diethyl phthallate, dibutylsebacate, and esters of citric acid among others. Generally, the plasticizers comprise between about 2% and about 50% by weight of polymer and plasticizer combined, preferably between about 5% and 15% by weight and most preferably about 10% by weight of the polymer and plasticizer combined.

The chewable tablets of this invention are prepared by spraying a solution of the methacrylic acid copolymer on to a fluidized bed of crystalline ibuprofen.

Crystalline ibuprofen can be prepared by the process described in U.S. Patent 4,476,248. The particle size of the ibuprofen should be about 80 to 500 microns and it it may contain excipients such as starch, lactose, hydroxypropyl methylcellulose, microcrystalline cellulose PVP, sucrose and fructose.

The residence time should be such that the ratio of copolymer to ibuprofen of each chewable tablet is about eight percent by weight.

The temperature of the inlet and outlet air should be maintained between 40 and 15 60 C. 20 and 40 C. respectively. The preferred inlet and outlet air temperatures are between 45 and 55 and 33 and 27°C respectively.

The temperature of the fluidized bed should be maintained between 60 and 20_p respectively. The preferred temperature of the bed is about 35 C.

The amount of ENDRAGIT L30D in the encapsulation formulation should be between about 10% to 60% by weight of ibuprofen, preferably about 14%.

15

EMBODIMENT OF THE INVENTION

Example 1 Preparation of chewable ibuprofen tablet (#13, p 23)

A. Encapsulated Ibuprofen

Ibuprofen 4 Kg
Eudragit L30D 2.33 Kg (coating polymer)
Propylene Glycol 140 gm (plasticizer)

Talc 200 gm Purified Water 200 gm

Ibuprofen crystals (particle size #40-105) are air suspended in a closed chamber (Glatt GPCG5). An aqueous dispersion of Eudragit L 30D and talc is sprayed onto the fluidized bed of ibuprofen at a rate of 60gm/min. The inlet and outlet air temperature are maintained at 50°C. and 20-22°C. respectively. The air rate is adjusted so as to maintain the particles in a suspended state and to maintain the fluidized bed at a temperature of 35°C.

Air Atomizing Pressure 3.5 bar

B. Preparation of Chewable Tablets

	•	Per Tab	Per 1000 Tab
	Mannitol	5-44 mg	544 Gm
	Malic Acid	4 mg	4 Gm
20	Aspartame	12 mg	12 Gm
	Spray Dried Orange Flavor	18 mg	18 Gm
	Encapsulated Ibuprofen	127 mg	127 Gm
•	Ac-Di-Sol	24 mg	24 Gm
25	Avicel pH102	60 mg	60 Gm
	F.D.C. Yellow 6 Lake	0.2 mg	0.2 Gm
	Citric Acid	4 mg	4 Gm
	Talc	20 mg	20 Gm

The compression mix of above was tabletted on a Manesty beta press with 1/2" flat face tooling. Tablet weight: 813 mg. Hardness: 9 - 13 Strong Cobert (SC). Disintegration Time in water: 2 minutes.

Table 1 shows a comparison of dissolution data between MOTRIN IB Tablet in PH 7.2 and the MOTRIN Chewable Tablets of this invention.

Table 2 shows a comparison of dissolution data between MOTRIN IB Tablets in PH 5.8 and the MOTRIN Chewable Tablets of this invention.

WO 91/15194 PCT/US91/01089

-7-

TABLE I

Comparison of Dissolution Data between Motrin IB
and Motrin Chewable Tablets

5 Buffer 7.2 PH

	Motri	n IB		#12		#13		rystallized
	•						Ibup	orofen Raw
				•			M	faterial
10							•	
	Flas			lask 2	Flas		Flas	
		% Released		% Released	Time	% Released		% Released
	0.10	-0.02	0.20	0.00	0.30	0.04	0.40	-0.02
	2.10	1.21	2.20	10.50	2.30	9.40	2.40	0.48
15	4.10	38.77	4.20	63.24	4.30	53.61	4.40	18.34
	6.10	71.49	6.20	89.31	6.30	84.02	6.40	40.63
	8.10	84.88	8.20	97.65	8.30	89.89	8.40	54.00
	10.10	90.60	10.20	95.85	10.30	90.91	10.40	63.24
	12.10	92.79	12.20	93.61	12.30	91.45	12.40	71.12
20	14.10	93.80	14.20	91.77	14.30	91.45	14.40	78.23
	16.10	94.36	16.20	91.38	16.30	91.62	16.40	85.25
	18.10	94.58	18.20	91.64	18.30	91.66	18.40	87.7 <i>5</i>
	20.10	94.67	20.20	91.86	20.30	91.68	20.40	94.43
	22.10	94.75	22.20	91.88	22.30	91.77	22.40	96.05
25	24.10	94.77	24.20	92.01	24.30	91.79	24.40	99.24
	26.10	94.82	26.20	92.16	26.30	91.77	26.40	100.63
	28.10	94.88	28.20	92.40	28.30	91.79	28.40	101.32
	30.10	94.92	30.20	92.57	30.30	91.81	30.40	101.58
	32.10	94.95	32.20	92.79	32.30	91.88	32.40	101.71
30	34.10	95.01	34.20	92.83	34.30	91.90	34.40	102.14
	36.10	94.99	36.20	92.92	36.30	91.86	36.40	102.20
	38.10	95.03	38.20	92.96	38.30	91.86	38.40	102.35
	40.10	94.99	40.20	93.00	40.30	91.84	40.40	102.46
	42.10	95.05	42.20	93.09	42.30	91.90	42.40	102.61
35	44.10	95.10	44.20	93.15	44.30	91.92	44.40	102.66
	46.10	95.08	46.20	93.20	46.30	91.97	46.40	102.72
	48.10	95.10	48.20	93.24	48.30	91.97	48.40	102.74
	50.10	95.08	50.20	93.26	50.30	91.94	50.40	102.76
	52.10	95.14	52.20	93.26	52.30	91.94	52.40	102.79
40	54.10	95.12	54.20	93.33	54.30	92.01	54.40	102.79
•	56.10	95.14	56.20	93.35	56.30	92.03	56.40	102.89
	58.10	95.18	58.20	93.37	58.30	92.05	58.40	102.85
	60.10	95.18	60.20	93.41	60.30	92.05	60.40	102.92

45 In PH7.2 buffer: Formula #12 shows fustic reline than Motrin IB.

Formula #12, #13 and Motrin IB shows the same reline forte after 10 minutes.

Formula #12, #13 all pass U.S.P. Tablets specification 20 minutes 785%.

Flask 1: Motrin IB.

Flask 2: Motrin chewable experiment Lot 12.

Flask 3: Motrin chewable experiment Lot 13.

Flask 4: Recrystallized Ibuprofen.

PCT/US91/01089

-9-

TABLE II

Buffer: 5-8 PH

5	Motrir	n IB	#12		#13		Recrysi Ibuproi Mater	en Raw
	Flas	sk 1	Flask 2		Flask 3		Flask 4	
10	Time 9	6 Released	Time	% Released	Time 9	% Released	Time %	Released
	0.10	0.04	0.20	0.02	0.30	0.00	0.40	0.04
	2.10	0.13	2.20	2.76	2.30	2.20	2.40	0.78
	4.10	2.25	4.20	9.78	4.30	7.90	4.40	9.18
	6.10	19.40	6.20	18.68	6.30	15.72	6.40	36.26
15	8.10	36.48	8.20	28.10	8.30	25.33	8.40	53.71
	10.10	48.08	10.20	36.05	10.30	34.69	10.40	65.90
	12.10	57.28	12.20	43.30	12.30	47.11	12.40	75.08
	14.10	64.34	14.20	51.49	14.30	56.54	14.40	82.38
	16.10	70.09	16.20	58.83	16.30	64.02	16.40	87.39
20	18.10	74.82	18.20	65.33	18.30	69.91	18.40	91.40
	20.10	78.57	20.20	71.02	20.30	74.86	20.40	94.41
	22.10	81.73	22.20	76.16	22.30	78.53	22.40	96.78
	24.10		24.20	80.43	24.30	81.51	24.40	98.47
	26.10	86.74	26.20	84.02	26.30	83.89	26.40	99.72
25	28.10	88.68	28.20	86.80	28.30	86.11	28.40	100.65
	30.10	90.24	30.20	89.05	30.30	87.78	30.40	101.51
	32.10	91.56	32.20	90.99	32.30	89.09	32.40	102.03
	34.10	92.70	34.20	92.53	34.30	90.06	34.40	102.53
	36.10	93.59	36.20	93.78	36.30	90.95	36.40	103.09
30	38.10	94.43	38.20	94.79	38.30	91.73	38.40	103.52
	40.10	95.21	40.20	95.62	40.30	92.46	40.40	103.91
	42.10	95.75	42.20	96.26	42.30	93.00	42.40	104.34
	44.10	96.29	44.20	96.76	44.30	93.50	44.40	104.67
	46.10	96.67	46.20	97.17	46.30	93.84	46.40	105.08
35	48.10	96.95	48.20	97.45	48.30	94.15	48.40	105.25
	50.10	97.21	50.20	97.65	50.30	94.36	50.40	105.46
	52.10	97.54	52.20	97.88	52.30	94.60	52.40	105.72
	54.10	97.75	54.20	98.12	54.30	94.77	54.40 56.40	105.94
40	56.10	97.97	56.20	98.25	56.30	94.97	56.40 58.40	106.11
40	58.10	98.12	58.20	98.38	58.30	95.08 05.18	58.40 60.40	106.29
	60.10	98.27	60.20	98.47	60.30	95.18	00.40	106.37

PH 5-8:

After 10 minutes about 10-12% difference action compare #12, #13 with Motrin IB.

After 20 minutes about 7-8% difference when compared #12, and #13 with Motrin IB.

After 30 minutes shows no significant difference among them.

-10-

Flask 1: Motrin IB.

Flask 2: Motrin chewable experiment Lot 12. Flask 3: Motrin chewable experiment Lot 13. Flask 4: Recrystallized Ibuprofen.

5

PCT/US91/01089

5

-11-

CLAIMS

- 1. A chewable taste-masked tablet having controlled release characteristics comprising a microcapsule of about 100 microns to about 0.8 mm in diameter having (a) a pharmaceutical core including crystalline ibuprofen and (b) a methacrylic acid copolymer coating having sufficient elasticity to withstand chewing.
- 2. A chewable taste-masked pharmaceutical composition according to claim 1 wherein the copolymer is Eudragit L30D said coating being adapted to release said 10 ibuprofen in the duodenum.
 - 3. The composition according to claim 1, wherein said copolymer coating further comprises a plasticizer.
- 15 4. The composition according to claim 2, wherein said copolymer coating further comprises a plasticizer.
- The composition according to claim 4, wherein said plasticizer is selected from
 the group consisting of glyceryl triacetate, triethyl citrate, acetyl triethyl citrate, dibutyl
 sebcate, acetyl tributyl citrate, diethyl phthlate, dibutyl phthlate, glycerine, propylene
 glycol and polyethylene glycol.
 - 6. The composition according to claim 5 wherein said plasticizer is propylene glycol.

25

INTERNATIONAL SEARCH REPORT

International Application No PCT/US 91/01089

I. CLASS	IFICATIO	N OF SUBJECT MATTER (if several classifi	ication symbols apply, indicate all) 6	
		ional Patent Classification (IPC) or to both Natio		
IPC ⁵ :	A 6	1 K 9/20, 9/50, 9/52,	31/19	
II. FIELDS	SEARCH	(ED		
		Minimum Documen	tation Searched 7	
Classification	on System	·	Classification Symbols	
IPC ⁵		A 61 K		
		Documentation Searched other to to the Extent that such Documents	nan Minimum Documentation are Included in the Fields Searched •	
III. DOCU	MENTS C	ONSIDERED TO BE RELEVANT		
Category •	Citat	ion of Document, 11 with Indication, where appr	opriate, of the relevant passages 12	Relevant to Claim No. 13
Y	EP,	A, 0212751 (FORMENTI) 4 March 1987 see claims 1-5,9,11,1 column 4, lines 33-47 lines 19-21 and 31; c	; column 5,	1-6
Y	EP,	A, 0212747 (PROCTER & 4 March 1987 see claims 1-3,6,7; p and 35; page 6, lines page 8, lines 28-29	eage 5, lines 24-29	1-6
Y	US,	A, 4835186 (AMERICAN 30 May 1989 see claims 1,2,5; col and 40		1-6
				
"A" doc con "E" earl filin "L" doc white cita "O" doc othe "P" doc late IV. CERT	ument defir sidered to I lier docume g date ument which ch is cited tion or othe ument refer er means ument publi r than the p	empletion of the international Search	"T" later document published after it or priority date and not in conflicited to understand the principle invention "X" document of particular relevant cannot be considered novel or involve an inventive step "Y" document of particular relevant cannot be considered to involve a document is combined with one ments, such combination being of in the art. "&" document member of the same published of Mailling of this international Section 177, 9	the with the application but or theory underlying the ce; the claimed invention cannot be considered to ce; the claimed invention an inventive step when the or more other such docubely to a person skilled catent family
22n	d May	1991) I, UI. 31	
Internation		eg Authority EAN PATENT OFFICE	Signature of Authorized Officer F.W. HECK	1000

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 9101089

SA 44948

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 24/06/91

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP-A- 0212751	04-03-87	JP-A- 6210301 US-A- 476601		
EP-A- 0212747	04-03-87	AU-B- 60169 AU-A- 611898 CA-A- 127504 GB-A,B 217925 JP-A- 6211192	6 19-02-87 8 09-10-90 4 04-03-87	
US-A- 4835186	30-05-89	None		

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82